

Evaluation of Physico-Chemical Characteristics in Groundwater Using GIS – A case Study of Chinnar Sub-basin, Cauvery River, Tamil Nadu, India

S.Venkateswaran¹, M.Elangomannan², M.Suresh¹, M.Vijay Prabhu¹,

1. Department of Geology, Periyar University, Salem – 11, Tamil Nadu.

2. Deputy Hydrogeologist, TWAD Board, Chennai – 16, Tamil Nadu.

provnkdswrn@gmail.com

Abstract

A detailed GIS based study on hydrochemistry of groundwater in Chinnar sub-basin, Dharmapuri District, Tamil Nadu, India has been Carried out to assess the quality of groundwater for determining its suitability for drinking purpose. Further, the spatial variation of various groundwater quality parameters over the basin has also been studied for November 2010. Fifty seven groundwater samples were collected and analysed for pH, conductance, total dissolved solids, total hardness, calcium, magnesium, fluoride, iron, carbonate, bicarbonate, total alkalinity, chloride, sodium, potassium, sulphate and nitrate etc., The values analyzed were evaluated in detail and compared with WHO water quality standards. TDS widely varied from 301 mg/L to 1505 mg/L with an average value of 703.51 mg/L. About 87% of the samples and spatially 789.32 km² areas are within the maximum allowable limit for drinking 1000 mg/L). Groundwater of the basin belongs to hard to very hard water category since the total hardness (TH) exceeds the permissible limit of 500 mg/L prescribed for drinking water. Magnesium content in groundwater fifty out of 57 samples exceeded the maximum allowable limit of 50 mg/L. Concentration of potassium ion in groundwater ranges from 3 mg/L to 24 mg/L with an average value of 9.11 mg/L. Thirty nine out of 57 samples exceeded the maximum allowable limit of 10 mg/L. Fluoride is (> 1.5 mg/L) in groundwater at 5 locations, which may cause dental and skeletal fluorosis.

Key words: Geographic Information System (GIS); World Health Organization (WHO); Spatial Variation; Dental and Skeletal Fluorosis.

Introduction

Water has a profound influence on human health and quality of the water supplied is important in determining the health of individuals and whole communities. Safe water quality is a major concern with reference to public health importance as health and well being of the human race is closely tied up with the quality of water used (Sharma et al. 2005). Despite major efforts to deliver safe piped, community water to the world's population, the

reality is that water supplies delivering safe water will not be available to all people in the near term (Agarwal 1981).

The quality of ground water depends on various chemical constituents and their concentration, which are mostly derived from the geological data of the particular region. Ground water occurs in weathered portion, along the joints and fractures of the rocks. In fact, industrial waste and the municipal solid waste have emerged as one of the leading cause of

pollution of surface and ground water. The principles governing the chemical characteristics of groundwater were well documented in many parts of the world (Garrels and Christ, 1965; Stumm and Morgan, 1970; Swaine and Schneider, 1971; Frapet al., 1984; Herczeg, et al., 1991; Som and Bhattacharya, 1992; Pawar, 1993; Wicks and Herman, 1994; Kimblin, 1995; Raju, 1998). This paper investigates the possible chemical processes of groundwater rock interaction in hard rock terrain.

GIS has emerged as a powerful technology for instruction, for research, and for building the stature of programs (Openshaw 1991; Longley 2000; Sui and Morrill 2004; Baker and Case 2000). Saraf et al., (1994) have conducted GIS based study and interpretation of groundwater quality data. Durbude et al., (2002) mapped the ground water quality parameters in Ghataprabha command area in GIS environment.

In the present study, groundwater samples have been collected and analyzed for various parameters such as, EC, pH, TDS, Ca, Mg, HCO₃, Cl, Na, K, Fe, F and NO₂ etc., the analysed results were taken in to GIS environment. In GIS, Spatial variation of groundwater quality parameters and their interrelationship have not been included. Further, it is observed that the concentration of major ions in groundwater of the area is high at many locations leading to unsuitability of groundwater for drinking. Thus, a GIS based study has been attempted to understand spatial variation of groundwater quality parameters over the Chinnar basin.

Study Area

The study area falls in Dharmapuri district of Tamil Nadu. Chinnar sub-basin, have been selected for the present investigation. It lies between 12°13'38" and 12°41'44" N latitudes, and 77°42'38" and 78°04'13" E longitudes covering an area of 893.65 Sq km out of which plain area covers 811.07 Sq km (Fig.1). Chinnar sub-basin is one of the major tributaries of Cauvery river. The basin comes under parts of Palakkode taluk and Pennagaram taluk of Dharmapuri district in Tamil Nadu State, India.

Methodology

57 groundwater samples from open and bore wells of various locations which are extensively used for drinking and also irrigation purposes in the Chinnar sub-basin area were collected during post-monsoon season (Nov. 2010). The locations of groundwater sampling stations are shown in the Fig. 1. Field parameters such as pH, electrical conductance (EC) and total dissolved solids (TDS) were measured immediately at sampling site using portable meters. Collected samples were brought to the laboratory on the same day, Ca and Mg were determined titrimetrically using standard EDTA, and chloride was determined by silver nitrate titration (Volgel, 1968). Carbonate and bicarbonate were estimated with standard sulphuric acid and sulphate was determined gravimetrically by precipitating BaSO₄ from BaCl₂. Na and K by Elico flame photometer (APHA, 1996).

The base map was prepared using toposheets on 1:50,000 scale. Their attributes are added and analyzed in ArcGIS software. Spatial analysis tools were used for the preparation of interpolation map. The maps were interpolated by using inverse distance methods to generate the spatial distribution map.

RESULTS AND DISCUSSION

Understanding the quality of groundwater is as important as that of its quantity, since, it is the main factor determining the suitability of water for drinking, domestic, agricultural and industrial purposes (Subramani & Elango 2005). The pH value is an important index of acidity or alkalinity and the concentration of hydrogen ion in groundwater (Murugesan et al. 2006). The lower value (pH < 4.0) will produce sour taste and higher value (pH > 8.5) an alkaline taste. The acceptable range of pH is normally 6.5 to 8.5 (WHO 1983). It is observed that the pH values of groundwater samples of the basin lie within the prescribed range showing an average value of 7.27.

The electrical conductivity (EC) is a measure of capability of water to transmit electrical current. It represents the total

concentration of soluble salts in water. It is used to measure the salinity hazard to crops as it reflects the TDS in groundwater (Anandakumar et al. 2007). The EC values in the study area vary widely from 430 $\mu\text{S}/\text{cm}$ to 2150 $\mu\text{S}/\text{cm}$ with an average value of 999.93 $\mu\text{S}/\text{cm}$. The higher values of EC may be due to long residence time and existing lithology of the region (Ballukraya & Ravi 1999).

Summary of the analytical results of various groundwater quality parameters is presented in Table 1. and the undesirable effects caused to humans when the parameters exceed the allowable limits (WHO 1983) are presented in Table 2. It is observed that potassium is 32% of samples present in exceeding permissible limit and TDS, Mg, K and Fe exceed the maximum allowable limits in more than 12% of the samples.

Total dissolved solids (TDS) range from 301 mg/L to 1505 mg/L with an average value of 703.51 mg/L. About 87% of the samples are within the maximum allowable limit for drinking (1000 mg/L) based on their TDS values (WHO 1983). The TDS spatial distribution map prepared using GIS (Fig. 2 and Table 5) reveals that groundwater in 21.74 km^2 of the area is unfit for drinking purposes. As per Freeze and Cherry (1979), 50 samples out of 57 represent the freshwater category (TDS < 1000 mg/L), and the remaining brackish water (TDS 1000 mg/L to 10000 mg/L) category (Table 3).

Total hardness (TH) also exhibits variation from 128 mg/L to 584 mg/L with an average value of 334.46 mg/L. Acceptable limit of TH for drinking is 500 mg/L (WHO 1983). The groundwater of the area is hard to very hard in nature because 54 samples (Table 4) have the TH values greater than 150 mg/L (Sawyer & McCarty 1967). The study area is delineated into three zones using GIS, based on the desirable (100 mg/L) and maximum permissible (500 mg/L) limits of TH as suggested by WHO (1983). The TH spatial distribution map (Fig. 3 and Table 5). illustrates that groundwater in 7.08 km^2 of the area is unsuitable for drinking purposes.

Magnesium content in groundwater of the area varies from 13 mg/L to 93 mg/L with an average value of 34.35 mg/L. Fifty out of 57 samples exceeded the maximum allowable limit of 50 mg/L for drinking as per the WHO (1996) standard. The Mg spatial distribution map (Fig. 4 and Table 5). illustrates that groundwater in 26.18 km^2 of the area is unsuitable for drinking purposes. Concentration of potassium ion in groundwater ranges from 3 mg/L to 24 mg/L with an average value of 9.11 mg/L. Thirty nine out of 57 samples exceeded the maximum allowable limit of 10 mg/L for drinking as per the WHO (1996) standard. The K spatial distribution map shows that groundwater quality based on K WHO limit 270.59 km^2 areas in the basin (Fig. 5 and Table 5).

Iron (Fe) content in groundwater of the area varies from 0 mg/L to 0.6 mg/L with an average value of 0.11 mg/L. Fifty out of 57 samples exceeded the maximum allowable limit of 0.3 mg/L for drinking as per the WHO (1983) standard. The Fe spatial distribution map (Fig. 6 and Table 5). reveals that 36.72 km^2 area is unsuitable for drinking purposes.

The chloride concentration varies from 24 mg/L to 308 mg/L. The average value is 98.11 mg/L. Fifty six samples exceeded the maximum allowable limit of 250 mg/L. The spatial distribution map indicates that 1.4 km^2 area is unsuitable for drinking purposes in the basin (Fig. 7 and Table 5).

Bedrock containing fluoride minerals is generally responsible for its high concentration in groundwater (Handa 1975, Wenzel & Blum 1992, Bardsen et al. 1996). The concentration of fluoride in groundwater of the basin varied from 0.46 mg/L to 2.3 mg/L with an average value of 1.04 mg/L. Ninety percent of the samples (5 out of 57) exhibited suitability for drinking purposes. The spatial distribution of fluoride concentration in groundwater during November 2010 is shows that 25.15 km^2 in Fig. 8 and Table 5.

Conclusions

The aforesaid statement reveals that the chemical composition of the Chinnar basin area

is hard, fresh to brackish, and slightly alkaline in nature. TDS about 87% of the samples and spatially 789.32 km² areas are within the maximum allowable limit for drinking 1000 mg/L). Total Hardness (TH) 7.08 km² area falls in exceeds the permissible limit of 500 mg/L prescribed for drinking water. Magnesium content in groundwater fifty out of 57 samples and 26.18 km² area fell in exceeded the maximum allowable limit of 50 mg/L. Concentration of potassium ion in groundwater ranges from 3 mg/L to 24 mg/L with an average value of 9.11 mg/L. Thirty nine out of 57 samples and 270.59 km² area fell in exceeded the maximum allowable limit of 10 mg/L. Fluoride is (> 1.5 mg/L) in groundwater at 5 locations and spatially 25.15 km² areas peoples affected for the dental and skeletal fluorosis in the Chinnar sub-basin this classification based on WHO standard for drinking purposes.

REFERENCES

- AGARWAL, A. 1981. Water, sanitation, health – for all. Center for international Development & Invironment, London 146 pp.
- ANANDAKUMAR, S., SUBRAMANI, T. AND ELANGO, L. 2007. Spatial variation of groundwater quality and inter elemental cOfTelation studies in Lower Bhavani River Basin, Tamil Nadu, India. Nat. Env. and Pollu. Tech., 6(2): 23.5-239.
- APHA (American Public Health Association) (1996) Standard methods for the Examination of water and wastewater 19th eds. Public Health Association, Washington, DC.
- BAKER THOMAS R., and CASE STEVEN B. (2000). Let GIS be your guide. The Science Teacher 67, no. 26. http://kangis.org/learning/publications/science_teacher/print/tst0010_24.pdf.
- BALLUKRAYA, P.N. AND RAVI, R.1999. Characteristic of groundwater in the unconfined aquifers of Chennai city. J. Geological Society of India, 54: 1-11.
- BARSEN, A., BJORVATN, K. AND SELVIG, KA. 1996. Variability in fluoride content of subsurface water reservoir,s. 1. Acta Odontol Scand.~ 54: 343-347.
- DURBUDE, D.G., VARADRAJAN, N., and PURANDARA, B.K. (2002) Mapping of ground water quality parameters in GIS environment, Proceeding of the International Conference on Hydrology and Water Management during 18-20 december, (2002) pp.568-577.
- FRAPE, S.K., FRITZ, P., and MCNUTT, R.H. (1984) Water rock interaction and chemistry of groundwaters from the Canadian Shield. Geochem. Cosmochim. Acta, v.48, pp.1617–1627.
- FREEZE, R.A. AND CHEERY, J.A. 1979. Text Book of Groundwater. Prentice Hall, Englewood Cliffs.
- GARRELS, R.M., and CHRIST, C.L. (1965) Solutions, Minerals and Equilibria. Harper and Row, New York, N.Y., 450p.
- HANDA, B..K. 1975. Geochemistry and genesis of fluoride containing groundwater in India Groundwater. J., 13(3)' 275-281.
- HERCZEG, A.L., TORGERSEN, T., CHIVAS, A.R., and HABERMEHL, M.A. (1991) Geochemistry of groundwater from the Great Artesian Basin, Australia. Jour. Hydrology, v.126, pp.225–245.
- KIMBLIN, R.T., (1995) The chemistry and origin of groundwater in Triassic sandstone and Quaternary deposits, Northwest England and some U.K. comparisons. Jour. Hydrology, v.172. pp.293–311.
- LONGLEY PAUL, A., (2000) The academic success of GIS in geography: Problems and prospects. Journal of Geographical Systems, 2 no. 1: pp.37 – 42.
- MURUGESAN, A., RAMU. A. AND KANNAN, N. 2006 Water quality assessment for Uttanlapalayam municipality in Theni District, Tamil Nadu, India. Poll. Res., 25(1): 163-166.
- OPENSHAW, S.A., (1991) view on the crisis in geography, or using GIS to put humpty-dumpty back together again. Environment and Planning, A 23, no. 5: pp.621-628.
- POORAN, N. J. (1993) Geochemistry of carbonate precipitation

- from the groundwaters in basaltic aquifers, An equilibrium thermodynamic approach, Jour. Geol. Soc. India, v.41, pp.119–131.
- RAJU, K.C.B. (1998) Importance of recharging depleted aquifers, State of the art of artificial recharge in India. Jour. Geol. Soc. India, v.51, pp.429–454.
- SARAF, A.K., GUPTA, R.P., JAIN, R.K., SWAIN, S., and SCHNEIDER, P. J., (1971) The chemistry of surface water in prairie ponds. Am. Chem. Soc. Adv. Chem. Ser., v.106, pp.99–104.
- SRIVASTAVA, N.K. (1994) GIS based processing and interpretation of ground water quality data, Proceedings of Regional workshop on Environmental Aspects of Ground water Development, Oct. 1994, Kurukshetra, India.
- SAWYER, C.N. AND MCCARTLY, D.L. 1967. Chemistry for Sanitary Engineers, 2nd edn. McGraw Hill, New York.
- SHARMA, S., R. BAJRACHARYA, B.K. SITAULA AND MERG . 2005. Water quality in the Central Himilaya, Current Science **89**(5):782.
- SOM. S.K., and BHATTACHARYA, A.K. (1992) Groundwater geochemistry of recent weathering at Panchpatmali bauxite bearing plateau, Koraput district, Orissa. Jour. Geol. Soc. India, v.40, pp.453–461.
- STUMM, W., and MORGAN, J.J. (1970) Aquatic Chemistry, Wiley, New York, N.Y. 1022p.
- SUBRALNANI, T. AND ELANGO, L. 2005. Groundwater quality and its suitability for drinking and agricultural use in Chithar river basin, Tamil Nadu, India. J. Env. Geo!,, 47: 1099-1110.
- SWAN DANIEL, and RICHARD MORRILL. (2004) Computers and geography: From automated geography to digital earth. In Geography and Technology, edited by STANLEY, D., BRUNN SUSAN, L., CUTTER, and J.W. HARRINGTON, JR. DORDRECHT, NL: KLUWER.
- WENZEL, W.W. AND BLUM, W.E.H. 1992. Fluoride speciation and mobility in fluoride contaminated soil and luinerals. J. Soil Sci., 153: 357-364.
- WHO, (1984) Guideline for drinking water quality. Vol.1. Recommendation, WHO, Genero, pp.1-4.
- WHO, (1996) Guideline for drinking water quality. Vol.2. Health criteria and other supporting information, WHO, Geneva. 973p.
- WIGGS, C.M., and HERMAN, J.S. (1994) The effect of a confining unit on the geochemical evolution of groundwater in the Upper Floridan aquifer system. Jour. Hydrology, v.153, pp.139–155.

Table 1. Chemic al Composition of Groundwater
(Ionic concentrations are expressed in mg/L and EC in μScm^{-1})

| Station | Ca | Mg | Na | K | Fe | HCO ₃ | CO ₃ | SO ₄ | Cl | NO ₃ | F | pH | EC* | TDS | TH |
|----------------|----|----|-----|----|----|------------------|-----------------|-----------------|-----|-----------------|-----|------|------|-----|-----|
| Achchattipalli | 90 | 40 | 104 | 12 | 0 | 344 | 0 | 64 | 108 | 17 | 1.2 | 6.58 | 1165 | 816 | 384 |
| Onnalvadi | 91 | 44 | 87 | 10 | 0 | 368 | 0 | 80 | 148 | 23 | 1.2 | 7.11 | 1273 | 891 | 412 |
| Oddarpalayam | 58 | 24 | 104 | 12 | 0 | 268 | 0 | 130 | 76 | 15 | 1.2 | 7.23 | 956 | 669 | 244 |

| | | | | | | | | | | | | | | | |
|-----------------------|-----|----|-----|----|-----|-----|---|-----|-----|----|-----|------|------|------|-----|
| Gulisandiram | 85 | 37 | 69 | 8 | 0 | 324 | 0 | 60 | 88 | 15 | 1.2 | 7.27 | 1009 | 706 | 368 |
| Bairamangalam | 86 | 39 | 63 | 8 | 0 | 336 | 0 | 78 | 96 | 17 | 1.2 | 7.04 | 1045 | 732 | 380 |
| Vanamangalam | 109 | 49 | 94 | 12 | 0.2 | 352 | 0 | 130 | 148 | 29 | 1.2 | 7.2 | 1372 | 960 | 476 |
| Masimayakkanappalli | 86 | 32 | 76 | 9 | 0.4 | 328 | 0 | 56 | 92 | 17 | 1.2 | 7.39 | 1013 | 709 | 376 |
| Karupalli | 91 | 42 | 112 | 14 | 0.6 | 388 | 0 | 78 | 128 | 29 | 1.2 | 7.28 | 1310 | 917 | 404 |
| Kelamangalam | 101 | 43 | 63 | 8 | 0 | 344 | 0 | 50 | 104 | 23 | 1.2 | 6.98 | 1093 | 965 | 340 |
| Belur | 114 | 49 | 74 | 9 | 0.1 | 396 | 0 | 80 | 156 | 29 | 2.1 | 7.41 | 1328 | 930 | 488 |
| Onnupalli | 104 | 40 | 124 | 14 | 0.3 | 388 | 0 | 95 | 136 | 32 | 1.2 | 7.36 | 1514 | 1060 | 448 |
| Adaikkalapuram | 104 | 40 | 118 | 14 | 0.2 | 396 | 0 | 74 | 132 | 27 | 0.7 | 7.44 | 1297 | 908 | 448 |
| Hosahalli | 131 | 58 | 178 | 24 | 0.4 | 612 | 0 | 110 | 208 | 27 | 1.6 | 7.21 | 1963 | 1374 | 568 |
| Anusonai | 37 | 14 | 36 | 5 | 0 | 168 | 0 | 27 | 32 | 3 | 0.7 | 7.45 | 444 | 311 | 152 |
| Ulimangalam | 72 | 32 | 65 | 8 | 0 | 260 | 0 | 110 | 80 | 15 | 1.2 | 7.25 | 949 | 664 | 312 |
| Modigapalayam | 42 | 19 | 52 | 6 | 0 | 208 | 0 | 21 | 48 | 6 | 0.7 | 6.62 | 574 | 402 | 184 |
| Makalgavundanu | 69 | 30 | 64 | 7 | 0 | 236 | 0 | 52 | 100 | 9 | 0.5 | 6.99 | 860 | 602 | 292 |
| Girisettipalli | 59 | 24 | 46 | 5 | 0 | 232 | 0 | 22 | 52 | 8 | 1.4 | 7.04 | 656 | 459 | 256 |
| Lakshampuram | 59 | 24 | 46 | 5 | 0 | 232 | 0 | 21 | 52 | 8 | 1.2 | 7.07 | 662 | 463 | 256 |
| Puram | 72 | 32 | 52 | 7 | 0 | 268 | 0 | 52 | 92 | 11 | 1.6 | 6.97 | 898 | 629 | 312 |
| Kammandur | 59 | 24 | 45 | 5 | 0 | 232 | 0 | 26 | 52 | 8 | 1.2 | 7.33 | 657 | 460 | 256 |
| Peddabaleguli | 134 | 60 | 158 | 22 | 0.3 | 532 | 0 | 120 | 244 | 38 | 1.2 | 6.95 | 1997 | 1398 | 584 |
| Basabanapalli | 35 | 14 | 43 | 6 | 0 | 188 | 0 | 25 | 36 | 6 | 1.2 | 7.46 | 552 | 386 | 164 |
| Doddakallupalli | 30 | 13 | 32 | 4 | 0 | 148 | 0 | 12 | 28 | 3 | 1.2 | 7.78 | 430 | 301 | 132 |
| Timijepalli | 72 | 29 | 36 | 5 | 0 | 264 | 0 | 38 | 48 | 8 | 0.5 | 7.05 | 744 | 521 | 300 |
| Odayandahalli | 88 | 36 | 72 | 9 | 0.3 | 316 | 0 | 50 | 112 | 17 | 0.5 | 7.42 | 1005 | 704 | 368 |
| Sandanapallu | 45 | 20 | 38 | 4 | 0 | 228 | 0 | 18 | 40 | 4 | 1.2 | 6.86 | 579 | 405 | 196 |
| Tippasandiram | 58 | 21 | 49 | 6 | 0.2 | 268 | 0 | 24 | 32 | 8 | 1.2 | 7.46 | 682 | 477 | 232 |
| Tattasandiram | 72 | 31 | 47 | 6 | 0 | 272 | 0 | 45 | 84 | 10 | 0.5 | 7.21 | 845 | 592 | 308 |
| Panchalli | 122 | 57 | 93 | 12 | 0.3 | 392 | 0 | 80 | 148 | 32 | 1.6 | 7.29 | 1480 | 1036 | 540 |
| Palaiyam | 106 | 45 | 78 | 9 | 0.2 | 408 | 0 | 95 | 104 | 29 | 0.5 | 6.89 | 1287 | 901 | 452 |
| Chudinur | 91 | 43 | 95 | 12 | 0.4 | 392 | 0 | 76 | 96 | 23 | 0.5 | 7.11 | 1259 | 881 | 436 |
| Unsatti | 29 | 13 | 48 | 5 | 0 | 180 | 0 | 20 | 28 | 4 | 0.5 | 7.33 | 486 | 340 | 152 |
| Nellukunti | 37 | 15 | 42 | 5 | 0 | 184 | 0 | 18 | 32 | 4 | 0.5 | 7.26 | 498 | 349 | 156 |
| Karadinattam | 106 | 45 | 79 | 10 | 0.3 | 408 | 0 | 72 | 136 | 29 | 1.2 | 7.89 | 1304 | 913 | 452 |
| Chinnabattagandahalli | 30 | 13 | 43 | 5 | 0 | 164 | 0 | 22 | 32 | 4 | 2.3 | 7.82 | 447 | 313 | 128 |
| Tottabadahalli | 45 | 20 | 52 | 7 | 0 | 228 | 0 | 30 | 44 | 6 | 1.2 | 7.42 | 608 | 426 | 196 |
| Agarharama | 122 | 57 | 93 | 12 | 0.4 | 392 | 0 | 80 | 148 | 42 | 1.2 | 7.39 | 1404 | 983 | 540 |
| Kombinaguddai | 101 | 44 | 66 | 8 | 0 | 320 | 0 | 60 | 124 | 27 | 1.2 | 7.31 | 1107 | 775 | 436 |
| Karadiguddai | 51 | 27 | 70 | 9 | 0.2 | 248 | 0 | 47 | 92 | 10 | 0.5 | 7.05 | 844 | 591 | 232 |
| Agaram | 86 | 42 | 94 | 12 | 0.4 | 352 | 0 | 76 | 116 | 21 | 1.2 | 7.51 | 1185 | 830 | 392 |
| Marandahalli | 107 | 52 | 112 | 16 | 0.2 | 376 | 0 | 85 | 204 | 36 | 1.2 | 7.43 | 1482 | 1037 | 484 |
| Bupanur | 45 | 20 | 58 | 7 | 0 | 216 | 0 | 21 | 72 | 5 | 1.2 | 7.35 | 671 | 470 | 196 |
| Gaalligattam | 32 | 14 | 27 | 3 | 0 | 172 | 0 | 11 | 28 | 4 | 0.5 | 7.72 | 449 | 314 | 140 |
| Sastramutlu | 67 | 29 | 98 | 12 | 0 | 312 | 0 | 54 | 112 | 19 | 1.2 | 7.36 | 1007 | 705 | 388 |
| Upparahalli | 102 | 49 | 85 | 10 | 0.4 | 428 | 0 | 80 | 124 | 27 | 1.2 | 6.88 | 1291 | 904 | 460 |
| Kesaraguli | 67 | 30 | 64 | 8 | 0 | 448 | 0 | 130 | 88 | 12 | 1.2 | 7.05 | 954 | 668 | 292 |
| Kariguddanur | 48 | 20 | 29 | 3 | 0 | 240 | 0 | 21 | 24 | 6 | 0.5 | 7.14 | 556 | 389 | 200 |
| Tirumalavadi | 42 | 18 | 53 | 6 | 0 | 196 | 0 | 23 | 68 | 5 | 1.2 | 7.38 | 636 | 445 | 180 |
| Attukottai | 70 | 33 | 43 | 5 | 0 | 272 | 0 | 60 | 92 | 8 | 1.2 | 7.43 | 883 | 618 | 312 |
| Eruduguttapatti | 59 | 27 | 40 | 5 | 0 | 248 | 0 | 90 | 44 | 6 | 1.2 | 7.69 | 763 | 534 | 260 |
| Budikal | 56 | 23 | 31 | 4 | 0 | 220 | 0 | 32 | 36 | 4 | 0.5 | 7.29 | 577 | 404 | 236 |
| Kadiyampatti | 74 | 34 | 30 | 4 | 0 | 284 | 0 | 43 | 48 | 40 | 1.2 | 7.32 | 801 | 561 | 324 |
| Siriyahalli | 128 | 56 | 192 | 24 | 0 | 436 | 0 | 85 | 308 | 44 | 1.2 | 7.65 | 2150 | 1505 | 532 |
| Ponnaiyankottai | 110 | 93 | 136 | 18 | 0.1 | 404 | 0 | 52 | 196 | 32 | 1.2 | 7.24 | 1520 | 1064 | 480 |
| Erranahalli | 86 | 36 | 93 | 12 | 0 | 340 | 0 | 48 | 140 | 23 | 0.9 | 7.56 | 1140 | 798 | 368 |
| Chakkilnattam | 112 | 43 | 90 | 12 | 0.1 | 372 | 0 | 64 | 156 | 27 | 1.2 | 7.26 | 1335 | 935 | 460 |

EC* – Electrical conductivity, RSC* – Residual Sodium Carbonate, SAR* – Sodium Adsorption Ratio, TH* - Total Hardness

Table 2. Groundwater samples of the study area exceeding the permissible limits prescribed by WHO standards for drinking purposes and the resulting undesirable effect on humans.

| Parameters | WHO International Standards 1983,1996) | | No. of samples Exceeding Permissible Limits | Total No. of Samples | Undesirable Effect on Human |
|--------------------------------------|--|--------------------------|--|----------------------|-----------------------------|
| | Most Desirable Limits | Maximum Allowable Limits | | | |
| pH | 6.5 - 8.5 | - | Nil | Nil | Taste |
| TDS (mg/l) | 500 | 1000 | 11,13,22,30,42,54,55. | 7 | Gastrointestinal Irritation |
| TH (mg/l) | 100 | 500 | 13,22,30,38,54. | 5 | Scale Formation |
| Ca ²⁺ (mg/l) | 75 | 200 | Nil | Nil | Scale Formation |
| Mg ²⁺ (mg/l) | 30 | 50 | 13,22,30,38,42,54,55. | 7 | Scale Formation |
| Na ⁺ (mg/l) | - | 200 | Nil | Nil | - |
| K ⁺ (mg/l) | - | 10 | 1,3,6,8,11,12,13,22,30,32,38,41,42,45,54,55,56,57. | 18 | - |
| Fe ²⁺ (mg/l) | - | 0.3 | 7,8,13,32,38,41,46. | 7 | Staining problem |
| Cl ⁻ (mg/l) | - | 250 | 54. | 1 | Salty taste |
| SO ₄ ²⁻ (mg/l) | - | 400 | Nil | Nil | Laxative effect |
| NO ₃ ⁻ (mg/l) | - | 45 | Nil | Nil | Blue baby disease |
| F ⁻ (mg/l) | - | 1.5 | 10,13,20,30,36. | 5 | Fluorosis |

Table 3. TDS Quality of groundwater based on Freeze and Cherry (1979),

| TDS)mg/L) | Nature of water | Nov. 2010 Representing Locations | Total No. Locations |
|----------------|-----------------|---|---------------------|
| < 1000 | Fresh water | 1,2,3,4,5,6,7,8,9,10,12,14,15,16,17,18,19,20,21,23,24,25,26,27,28,29,31,32,33,34,35,36,37,38,39,40,41,43,44,45,46,47,48,49,50,51,52,53,56,57. | 50 |
| 1000 - 10000 | Brackish water | 11,13,22,30,42,54,55. | 7 |
| 10000 - 100000 | Saline water | - | - |
| > 100000 | Brine water | - | - |

Table 4. Classification of groundwater based on hardness

| Total Hardness as CaCO ₃ (mg/l) | Water Class | Representing Locations Nov. 2010 | Total No. of Locations |
|--|-----------------|--|------------------------|
| < 75 | Soft | Nil | Nil |
| 75 – 150 | Moderately hard | 24,36,44. | 3 |
| 150 – 300 | Hard | 3,14,16,17,18,19,21,23,25,27,28,33,34,37,40,43,47,48,49,51,52. | 21 |

| | | | |
|-------|-----------|---|----|
| > 300 | Very hard | 1,2,4,5,6,7,8,9,10,11,12,13,15,20,22,26,29,30,31,32,35,38,39,41,42,45,46,50,53,54,55,56,57. | 33 |
|-------|-----------|---|----|

Table 5. Chemical Quality – GIS Spatial Distribution Results

| Class | TDS – Results Area in km ² | TH – Results Area in km ² | Mg – Results Area in km ² | K – Results Area in km ² | Cl – Results Area in km ² | Fe – Results Area in km ² | F – Results Area in km ² |
|--------------------------|---------------------------------------|--------------------------------------|--------------------------------------|-------------------------------------|--------------------------------------|--------------------------------------|-------------------------------------|
| Most Desirable Limits | 98.54 | 0.07 | 284.55 | - | - | - | - |
| Maximum Allowable Limits | 690.78 | 803.93 | 500.34 | 540.47 | 809.59 | 774.35 | 785.92 |
| Exceeding Permissible | 21.74 | 7.08 | 26.18 | 270.59 | 1.48 | 36.72 | 25.15 |




Fig. 1. Study Area And Water Sample Location Map




Fig. 2. TDS Spatial Distribution Map




Fig. 3. TH Spatial Distribution Map




Fig. 4. Mg Spatial Distribution Map




Fig. 5. K Spatial Distribution Map




Fig. 6. Cl Spatial Distribution Map




Fig. 7. Fe Spatial Distribution Map




Fig. 8. F Spatial Distribution Map